

TCCC Journal Watch

August 2017

Abdallah N, Salama A, Elithy A: Effects of preincisional analgesia with surgical site infiltration of ketamine or levobupivacaine in patients undergoing abdominal hysterectomy under general anesthesia; a randomized double blind study. *Saudi J Anaesth* 2017;11:267-272

Abdolghaffari A, Panahi Y, Sahebkar A: A review on symptoms, treatments protocols and proteomic profile in sulfur mustard-exposed victims. *J Cell Biochem* 2017;Epub ahead of print

Adamiak Z, Krystkiewicz W, Pomianowski A, et al: The effect of hemostatic dressing prototypes for the uniformed services on selected blood coagulation parameters in pigs. *Acta Vet Scand* 2017;59:29

Alkhouri H, Vassiliadis J, Murray M, et al: Emergency airway management in Australian and New Zealand emergency departments: a multicenter descriptive study of 3710 emergency intubations. *Emerg Med Australas* 2017;Epub ahead of print

Amer K, Rehman S, Haydel C: Efficacy and safety of tranexamic acid in orthopedic fracture surgery: a meta-analysis and systematic literature review. *J Orthop Trauma* 2017;Epub ahead of print

Amir S, Maya S, Irina R, et al: Pre hospital administration of freeze dried plasma, is it the solution for trauma casualties? *J Trauma Acute Care Surg* 2017;Epub ahead of print

Ariffin A: Tube thoracostomy complications: more to learn. *World J Surg* 2017;Epub ahead of print

Asehnoune K, Balogh Z, Citerio G, et al: The research agenda for trauma critical care. *Intensive Care Med* 2017;Epub ahead of print

Aykanat Ö, Karakoyun D, Türkoğlu M, Dinc C: Anti-edematous, anti-inflammatory and neuroprotective effect of etanercept in acute stage in experimental head injury. *Ulus Travma Acil Cerrahi Derg* 2017;23:173-180

Baatista A, Wright T, Meany J, et al: Red cell storage duration does not affect outcome after massive blood transfusion in trauma and nontrauma patients: a retrospective analysis of 305 patients. *Biomed Res Int* 2017;Epub ahead of print

Badenes R, Oddo M, Suarez J, et al: Hemoglobin concentrations and RBC transfusion thresholds in patients with acute brain injury: an international survey. Crit Care 2017;21:159

Barbier O, Malgras B, Choufani C, et al: Surgical support during the terrorist attacks in Paris, November 13, 2015: experience at Bégin military teaching hospital. J Trauma Acute Care Surg 2017;82:1122-1128

Barnard E, Manning J, Smith J, et al: A comparison of selective aortic arch perfusion and resuscitative endovascular balloon occlusion of the aorta for the management of hemorrhage-induced traumatic cardiac arrest: a transitional model in large swine. PLoS Med 2017;14:e1002349

Barron M, Kuckelman J, McClellan J, et al: Smartphone-based mobile thermal imaging technology to assess limb perfusion and tourniquet effectiveness under normal and blackout conditions. J Trauma Acute Care Surg 2017;Epub ahead of print

Bayer J, Lefering R, Reinhardt S, et al: Severity-dependent differences in early management of thoracic trauma in severely injured patients- analysis based on the TraumaRegister DGU. Scand J Trauma Resusc Emerg Med 2017;25:10

Beavin A, Briard R, Ballard M, Parker P: Two new effective tourniquets for potential use in the military environment: a serving soldier study. Mil Med 2017;182:e1929-e1932

Belin T, Yazer M, Meledeo M, et al: An evaluation of methods of producing low-titer group O whole blood to support military trauma resuscitation. J Trauma Acute Care Surg 2017;82:S79-S86

Bennett B, Holcomb J: Battlefield trauma-induced hypothermia: transitioning the preferred method of casualty rewarming. Wilderness Environ Med 2017;Epub ahead of print

Benov A, Salas M, Nakar H, et al: Battlefield pain management: a view of 17 years in Israel Defense Force. J Trauma Acute Care Surg 2017;83:S150-S155

Benov A, Yaslowitz O, Hakim T, et al: The effect of blood transfusion on compensatory reserve: a prospective clinical trial. J Trauma Acute Care Surg 2017;83:S71-S76

Beranger F, Lesquen H, Aoun O, et al: Management of war-related vascular wounds in French Role 3 hospital during the Afghan campaign. Injury 2017;Epub ahead of print

Berséus O: Effects on the anti-ABO titers of military blood donors from a predeployment vaccination program. J Trauma Acute Care Surg 2017;82:S91-S95

Bilge S, Aydin A, Bilge M, et al: A study on the tactical safety of endotracheal intubation under darkness. Mil Med 2017;182:e1722-e1725

Bordes J, Joubert C, Esnault P, et al: Coagulopathy and transfusion requirements in war related penetrating traumatic brain injury. A single centre study in French role 3 medical treatment facility in Afghanistan. Injury 2017;48:1047-1053

Bowman J, Juergens A, McClure M, Spear D: Intubation of the right atrium during an attempted modified surgical airway in a pig. J Spec Oper Med 2017;17:96-100

Bunya N, Harada K, Kuroda Y, et al: The effectiveness of hybrid treatment for severe multiple trauma: a case of multiple trauma for damage control laparotomy and thoracic endovascular repair. Int J Emerg Med 2017;10:18

Butler F: Fluid resuscitation in Tactical Combat Casualty Care: yesterday and today. Wilderness Environ Med 2017;28:S74-S81

Bynum J, Taylor A, Peltier G, et al: Evaluation of adenosine, lidocaine, and magnesium for enhancement of platelet function during storage. J Trauma Acute Care Surg 2017;83:S9-S15

Callaway D, Puciaty A, Robertson J, et al: Care Report: life saving application of commercial tourniquet in pediatric extremity hemorrhage. Prehosp Emerg Care 2017;28:1-3

Cannon J, Johnson M, Caskey R, et al: High ratio plasma resuscitation does not improve survival in pediatric trauma patients. J Trauma Acute Care Surg 2017;83:211-217

Carness J, Wilson M, Lenart M, et al: Experience with regional anesthesia for analgesia during prolonged aeromedical evacuation. Aerosp Med Hum Perform 20017;88:768-772

Chandler H, MacLeod K, Penn-Barwell J, et al: Extremity injuries sustained by the UK military in the Iraq and Afghanistan conflicts: 2003-2014

Chang R, Eastridge B, Holcomb J: Remote damage control resuscitation in austere environments. Wilderness Environ Med 2017;28:S124-S134

Chang R, Fox E, Greene T, et al: Multicenter retrospective study on noncompressible torso hemorrhage: anatomic locations of bleeding and comparison of endovascular versus open approach. *J Trauma Acute Care Surg* 2017;83:11-18

Chang R, Holcomb J, Johansson P, et al: Plasma resuscitation improvised survival in a cecal ligation and puncture rat model of sepsis. *Shock* 2014;Epub ahead of print

Chen W, Van Buren P: A case of severe neutropenia from short-term exposure to moxifloxacin. *J Investig Med High Impact Case Rep* 2017;Epub ahead of print

Christie A, Kornblith L, Howard B, et al: Characterization of distinct coagulopathic phenotypes in injury: pathways-specific drivers and implications for individualized treatment. *J Trauma Acute Surg* 2017;82:1055-1062

Convertino V, Sawaka M: Wearable technology for compensatory reserve to sense hypovolemia. *J Appl Physiol* 2017;Epub ahead of print

Convertino V, Schiller A: Measuring the compensatory reserve to identify shock. *J Trauma Acute Care Surg* 2017;82:S57-S65

Cornelius B, Campbell R, McGauly P: Tourniquets in trauma care: a review of application. *J Trauma Nurs* 2017;24:203-207

Corso P, Mandry C, Reynolds S: The sole provider: preparation for deployment to a medically austere theater. *J Spec Oper Med* 2017;17:74-81

Costantini T, Coimbra R, Holcomb J, et al: Pelvic fracture pattern predicts the need for hemorrhage control intervention- results of an AAST multi-institutional study. *J Trauma Acute Care Surg* 2017;82:1030-1038

Daniel Y, Sailliol A, Pouget T, et al: Whole blood transfusion closest to the point-of-injury during French remote military operations. *J Trauma Acute Care Surg* 2017;82:1138-1146

David J, Bouzat P: Early fibrinogen-concentrate administration in management of trauma-induced coagulopathy. *Lancet Haematol* 2017;4:e348

David J, Voiglio E, Cesareo E, et al: Prehospital parameters can help to predict coagulopathy and massive transfusion in trauma patients. *Vox Sang* 2017;Epub ahead of print

Davidson A, Russo R, Ferenez S, et al: Incremental balloon deflation following complete resuscitative endovascular balloon occlusion of the aorta results in steep inflection of flow and rapid reperfusion in a large animal model of hemorrhagic shock. *J Trauma Acute Care Surg* 2017;83:139-143

Dekker S, Nokolian V, Sillesen M, et al: Different resuscitation strategies and novel pharmacologic treatment with valproic acid in traumatic brain injury. *J Neurosci Res* 2017;Epub ahead of print

Denninghoff K, Nuño T, Pauls Q, et al: Prehospital intubation is associated with favorable outcomes and lower mortality in ProTECT III. *Prehosp Emerg Care* 2017;Epub ahead of print

Dents C, Bradley M, Schobel S, et al: Towards precision medicine: accurate predictive modeling of infectious complications in combat casualties. *J Trauma Acute Care Surg* 2017;Epub ahead of print

Dhingra D, Singh A, Masood O: Safety pin as a chest tube clamp - cause of iatrogenic pneumothorax. *Indian J Pediatr* 2017;Epub ahead of print

Diebel L, Martin J, Liberati D: Early tranexamic acid administration ameliorates the endotheliopathy of trauma and shock in an in vitro model. *J Trauma Acute Care Surg* 2017;82:1080-1086

Dilmen S, Eryilmaz M, Balkan S, et al: Comparison of warm fluid and cold fluid resuscitation during uncontrolled hemorrhagic shock model in rats. *Ulus Travma Acil Cerrahi Derg* 2017;23:287-293

Drake S, Wolf D, Meininger J, et al: Methodology to reliably measure preventable trauma death rate. *J Trauma Surg Acute Care* 2017;Epub ahead of print

Driver B, Klein L, Schick A, et al: The occurrence of aspiration pneumonia after emergency endotracheal intubation. *Am J Emerg Med* 2017;Epub ahead of print

Dubose J: How I do it: partial resuscitative endovascular balloon occlusion of the aorta (P-REBOA). *J Trauma Acute Care Surg* 2017;83:197-199

Duggan L, Lockhart S, Romano K, et al: Front-of-neck airway meets front-of-neck simulation: improving cricothyroidotomy skills using a novel open-access three-dimensional model and the airway app. *Can J Anaesth* 2017;Epub ahead of print

Dunbar N, Yazer M, Biomedical Excellence for safer Transfusion (BEST) Collaborative and the STAT Study Investigators. *Transfusion* 2017;Epub ahead of print

Ekaney M, Grable M, Powers W, et al: Cytochrome c and resveratrol preserve platelet function during cold storage. *J Trauma Acute Care Surg* 2017;83:271-277

Feng J, Wang W, Zeng J, et al: Optimization of brain metabolism using metabolic-targeted therapeutic hypothermia can reduce mortality from traumatic brain injury. *J Trauma Acute Care Surg* 2017;Epub ahead of print

Fisher A, Teeter W, Cordova C, et al: The role I resuscitation team and resuscitative endovascular balloon occlusion of the aorta. *J Spec Oper Med* 2017;17:65-73

Fitzgerald M, Esser M, Russ M, et al: Pelvic trauma mortality reduced by integrated trauma care. *Emerg Med Australas* 2017;Epub ahead of print

Fouche P, Jennings P, Smith K, et al: Survival in out-of-hospital rapid sequence intubation for non-traumatic brain injury. *Prehosp Emerg Care* 2017;Epub ahead of print.

Frank B, Maher Z, Hazelton J, et al: Venous thromboembolism after major venous injuries: competing priorities. *J Trauma Acute Care Surg* 2017;Epub ahead of print

Gaither J, Ckani V, Stoltz U, et al: Body temperature after EMS transport: association with traumatic brain injury outcomes. *Prehosp Emerg Care* 2017;Epub ahead of print

Galante J: Using tourniquets to stop bleeding. *JAMA* 2017;317:1490

Galvagno S, Fox E, Appana S, et al: Outcomes following concomitant traumatic brain injury and hemorrhagic shock: a secondary analysis for the PROPPR Trial. *J Trauma Acute Care Surg* 2017;Epub ahead of print

Georgeff P, Nikolian V, Halaweish I, et al: Resuscitation with lyophilized plasma is safe and improves neurological recovery in long-term survival model of swine subjected to traumatic brain injury, hemorrhagic shock, and polytrauma. *J Neurotrauma* 2017;34:2167-2175

Gonzalez-Cortes J, Bages-Rousselon Y, Gonzalez-Cantu J, Mohamed-Hamsho J: Minimally invasive surgery for the removal of posterior intraocular foreign bodies. *J Ophthalmic Vis Res* 2017;12:236-240

Gooble S, Frank S: Tranexamic acid: What is known and unknown, and where do we do from here? Anesthesiology

Grocott H: Definitive airway management in the presence of a laryngeal tube supraglottic airway: "There's more than one way to skin a cat". A Case Rep 2017;Epub ahead of print

Gurney J, Holcomb J: Blood transfusion form the military's standpoint: making last century's standard possible today. Curr Trauma Rep 2017;3:144-155

Haghghi M, Ettehad H, Mardani-Kivi M, et al: Does tranexamic acid reduce bleeding during femoral fracture operation? Arch Bone Jt Surg 2017;5:103-108

Hartch P, Pogarty E: Subacute normobaric oxygen and hyperbaric oxygen therapy in drowning, reversal of brain volume loss: a case report. Med Gas Res 2017;7:144-149

Harvey M, Hunt R, Hick J, Hanfling D: The role that health care coalitions play in support of mass casualty response. Ann Emerg Med 2017;70:106-107

Harvin J, Maxim T, Inaba K, et al: Mortality following emergent trauma laparotomy: a multicenter, retrospective study. J Trauma Acute Care Surg 2017;Epub ahead of print

Henriksen H, Grand A, Viggers S, et al: Impact of blood products on platelet function in patients with traumatic brain injuries: a translation study. J Surg Res 2017;14:154-161

Hernandez M, Thiels C, Aho J, et al: Prehospital plasma resuscitation is associated with improved neurologic outcomes after traumatic brain injury. J Trauma Acute Care Surg 2017;Epub ahead of print

Hill K: Association of bystander cardiopulmonary resuscitation with overall & neurologically favorable survival after pediatric out-of-hospital cardiac arrest in the United States. J Emerg Med 2017;52:905

Hinson H, Rowell S, Morris C, et al: Early fever after trauma: does it matter? J Trauma Acute Care Surg 2017;Epub ahead of print

Holcomb J: Major scientific lessons learned in the trauma field over the last two decades. PLoS Med 2017;14:e1002339

Holcomb J, Swartz M, DeSantis S, et al: Multicenter observational prehospital resuscitation on helicopter study. J Trauma Acute Care Surg 2017;S83-S91

Horn G, Bowling F, Lowe D, et al: Manikin human-patient simulator training. *J Spec Oper Med* 2017;17:89-95

Howard J, Stockinger Z, Cap A, et al: Military use of TXA in combat trauma: does it matter? *J Trauma Acute Care Surg* 2017;Epub ahead of print

Huang G, Dunahm C: Mortality outcomes in trauma patients undergoing prehospital red blood cell transfusion: a systematic literature review. *Int J Burns Trauma* 2017;7:17-26

Huebner B, Moore E, Moorre H, et al: Freeze-dried plasma enhance clot formation and inhibits fibrinolysis in the presence of tissue plasminogen activator similar to pooled liquid plasma. *Transfusion* 2017;Epub ahead of print

Huff C: Nonclinicians get crash course in bleeding control. *Ann Emerg Med* 2017;69:18A-20A

Hydmo P, Horodyski M, Conrad B, et al: Does the novel lateral trauma position cause more motion in an unstable cervical spine injury than the logroll maneuver? *Am J Emerg Med* 2017;Epub ahead of print

Iapichino G, Ponschab M, Cadamuro J, et al: Concentrated lyophilized plasma used for reconstitution of whole blood leads to higher coagulation factor activity but unchanged thrombin potential compared with fresh-frozen plasma. *Transfusion* 2017;57:1763-1771

Innerhofer P, Fries D, Mittermayr M, et al: Reversal of trauma-induced coagulopathy using first-line coagulation factor concentrates or fresh frozen plasma (RETIC): a single-centre, parallel-group, open-label, randomised trial. *Lancet Haematol* 2017;Epub ahead of print

Innerhofer P, Fries D, Oswald E, RETIC Study Group: Early fibrinogen-concentrate administration in management of trauma induced coagulopathy-author's reply. *Lancet Haematol* 2017;4:e348-e349

Inocencio M, Childs J, Chilstrom M, Berona K: Ultrasound findings in tension pneumothorax: a case report. *J Emerg Med* 2017;52:e217-e220

Jacobs B, Cain-Nielsen A, Jakubus J, et al: Unfractionated heparin versus low-molecular-weight heparin for venous thromboembolism prophylaxis in trauma. *J Trauma Acute Care Surg* 2017;83:151-158

Jaffee E, Alpert E, Lipsky A: A unique program to incorporate volunteers into a nationwide emergency medical system: maximizing preparedness for a mass casualty incident. *JAMA Surg* 2017;Epub ahead of print

Jarman A, Hopkins C, Hasen J, et al: Advanced airway type and its association with chest compression interruptions during out-of-hospital cardiac arrest resuscitation. *Prehosp Emerg Care* 2017;Epub ahead of print

Jodie P, Kerstin H: BET 1: Pre-hospital finger thoracostomy in patients with chest trauma. *Emerg Med J* 2017;34:417-418

Johnson M, Alarhayem A, Convertino V, et al: Compensatory reserve index: performance of a novel monitoring technology to identify the bleeding trauma patient. *Shock* 2017;Epub ahead of print

Johnson M, Williams T, Ferenez S, et al: The effect of resuscitative endovascular balloon occlusion of the aorta, partial aortic occlusion and aggressive blood transfusion on traumatic brain injury in a swine with multiple injuries model. *J Trauma Acute Care Surg* 20017;83:61-70

Jones C, Rothermel A, Mackay D: Evidence-based medicine: wound management. *Plast Reconstr Surg* 2017;140:201e-216e

Joshi N, Hayes B, Mason J, Grock A: Does the intranasal route pass the sniff test? *Ann Emerg Med* 2017;70:212-214

Kaji A, Bosson N, Gausche-Hill M, et al: Patient outcomes in urban and suburban level I versus level II trauma centers. *Ann Emerg Med* 2017;70:161-168

Keenan S, Riesberg J: Prolonged field care: beyond the “golden hour”. *Wilderness Environ Med* 2017;28:S135-S139

Khan M, Streets C, Tai N, Rickard R: United Kingdom military surgical preparedness for contingency operations. *J Trauma Acute Care Surg* 2017;83:S142-S144

Kheirabadi B, Terrazas I, Miranda N, et al: Do vented chest seals differ in efficacy? An experimental evaluation using a swine hemopneumothorax model. *J Trauma Acute Care Surg* 2017;83:182-189

King B, Peterson W: The care of thermally injured patients in operational, austere and mass casualty situations. *Wilderness Environ Med* 2017;28:S103-S108

Kirkpatrick A, McKee J, McBeth P, et al: The damage control surgery in austere environment research group (DCSAERG): a dynamic program to facilitate real-time telementoring/telediagnosis to address exsanguination in extreme in and austere environments. *J Trauma Acute Care Surg* 2017;83:S156-S163

Klebacher R, Harris M, Ariyaprakai N, et al: Incidence of naloxone redosing in the age of the new opioid epidemic. *Prehosp Emerg Care* 2017;Epub ahead of print

Klein L, Huelster J, Adil U, et al: Endovascular rewarming in the emergency department for moderate to severe accidental hypothermia. *Am J Emerg Med* 2017;Epub ahead of print

Koko K, McCauley B, Gaughan J, et al: Kaolin-based hemostatic dressing improves hemorrhage control from a penetrating inferior vena cava injury in coagulopathic swine. *J Trauma Acute Care Surg* 2017;83:71-76

Kotwal R, Montgomery H, Miles E, et al: Leadership and a casualty response system for eliminating preventable death. *J Trauma Acute Care Surg* 2017;82:S9-S15

Kragh J, Aden J, Shackelford S, et al: Assessment of trainer skill to control groin-wound bleeding: use of junctional tourniquet models on a manikin. *J Spec Oper Med* 2017;17:39-48

Kragh J, Dubick M: Bleeding control with limb tourniquet use in the wilderness setting: review of science. *Wilderness Environ Med* 2017;28:S25-S32

Kroll M, Ritter M, Williams H: Fatal and non-fatal burn injuries with electrical weapons and explosive fumes. *J Forensic Leg Med* 2017;50:6-11

Kronen R, Liang S, Bochicchio G, et al: Invasive fungal infections secondary to traumatic injury. *Int J Infect Dis*;2017;Epub ahead of print

Kumar K, Kumar J, Sarvagna J, et al: Hemostasis and post-operative care of oral surgical wounds by Hemcon dental dressing in patients on oral anticoagulant therapy: a split mouth randomized controlled clinical trial. *J Clin Diagn Res* 2016;10:ZC37-ZC40

Lallemand M, Moe, D, McClellan J, et al: Resuscitative endovascular balloon occlusion of the aorta for major abdominal venous injury in a porcine hemorrhagic shock model. *J Trauma Acute Care Surg* 2017;83:230-236

Landsdal H, Berge M, Kristensen F, et al: A strategy for securing a definitive airway after successful rescue by supraglottic airway device: TABASCO. *Acta Anaesthesiol Scand* 2017;Epub ahead of print

Leathermann M, Held J, Fluke L, et al: Relative device stability of anterior versus axillary needle decompression for tension pneumothorax during casualty movement: preliminary analysis of a human cadaver model. *J Trauma Acute Care Surg* 2017;83:S136-S141

Lee Y, Huang Y, Hung M, et al: Risk factors associated with the development of seizures among adult patients treated with ertapenem: a matched case-control study. *PLoS One* 2017;12:e0182046

Lesaffre X, Tourtier J, Violin Y, et al: Remote damage control during the attacks of Paris: lessons learned for the Paris Fire Brigade and evolutions in the rescue system. *J Trauma Acute Care Surg* 2017;82:S107-S113

Letson H, Dobson G: 3% NaCl adenosine, lidocaine, Mg²⁺ (ALM) bolus and 4 hours "drip" infusion reduces noncompressible hemorrhage by 60% in a rat model. *J Trauma Acute Care Surg* 2017;82:1063-1072

Littlejohn L: Treatment of thoracic trauma: lessons from the battlefield adapted to all austere environments. *Wilderness Environ Med* 2017;28:S69-S73

Liu N, Holcomb J, Wade C, et al: Inefficacy of standard vital signs for predicting mortality and the need for prehospital life-saving interventions in blunt trauma patients transported via helicopter: a repeated call for new measures. *J Trauma Acute Care Surg* 2017;83:S98-S103

Liu Y, Li S, Li Z, et al: A safety evaluation of profound hypothermia-induced suspended animation for delayed resuscitation at 90 or 120 min. *Mil Med Red* 2017;4:16

Llewellyn C: The symbiotic relationship between operational military medicine, tactical medicine, and wilderness medicine: a view through a personal lens. *Wilderness Environ Med* 2017;28:S6-S11

Lloyd B, Murray C, Shaikh F, et al: Early infectious outcomes following addition of fluoroquinolone or aminoglycoside to post-trauma antibiotic prophylaxis in combat-related open fractures injuries. *J Trauma Acute Care Surg* 2017; Epub ahead of print

Long B, Koyfam A: Best clinical practice: controversies in outpatient management of acute pulmonary embolism. *J Emerg Med* 20017;52:668-679

Louro J, Anderson K, Dudaryk R: Correction of severe coagulopathy and hyperfibrinolysis by tranexamic acid and recombinant Factor VIIa in a cirrhotic patient after trauma: a case report. *A A Case Rep* 2017;Epub ahead of print

Mabry R, Kharod C, Bennett B: Awake cricothyrotomy: a novel approach to the surgical airway in the tactical setting. *Wilderness Environ Med* 2017;28:S61-S68

Mackenzie C, Pasley J, Garofalo E, et al: Head-camera video recordings of trauma core competency procedures can evaluate surgical resident's technical performance as well as colocated evaluators. *J Trauma Acute Care Surg* 2017;83:S124-S129

Maddry J, Mora A, Savell S, et al: Impact of critical care air transport team (CCATT) ventilator management on combat mortality. *J Trauma Acute Care Surg* 2017;Epub ahead of print

Mahmood A, Roberts I, Shakur H: A nested mechanistic sub-study into the effect of tranexamic acid versus placebo on intracranial haemorrhage and cerebral ischaemia in isolated traumatic brain injury: study protocol for a randomised controlled trial (CRASH-3 trial intracranial bleeding mechanistic sub-study [CRASH-3 IBMS]). *Trials* 2017;18:330

Maissan I, Ketelaars R, Vlottes B, et al: Increase in intracranial pressure by application of a rigid cervical collar: a pilot study in healthy volunteers. *Eur J Emerg Med* 2017;Epub ahead of print

Malhotra R, Chua W, O'Neill G: Calf compartment syndrome associated with the use of an intraosseous line in an adult patient: a case report. *Malays Orthop J* 2016;10:49-51

Marehbian J, Muehlschlegel S, Edlow B, et al: Medical management of the severe traumatic brain injury patient. *Neurocrit Care* 2017;Epub ahead of print

McDaniel S, Golla S, Moore A, et al: Resuscitation of hypotensive traumatic brain injured animals with spray-dried plasma does not adversely affect physiology and improves blood-brain barrier function. *Mil Med* 2017;182:e1706-e1711

McQuilten Z, Bailey M, Cameron P, et al: Fibrinogen concentration and use of fibrinogen supplementation with cryoprecipitate in patients with critical bleeding

receiving massive transfusion: a bi-national cohort study. Br J Haematol 2017;Epub ahead of print

Mead K, Tennent D, Stinner D: The importance of medical readiness training exercise: maintaining medical readiness in a low-volume combat casualty flow era. Mil Med 2017;182:e1734-e1737

Meledeo M, Fisher A, Peltier G, et al: Volumetric control of whole blood collection in austere environments. J Trauma Acute Care Surg 2017;82:S26-S32

Meledeo M, Herzig M, Bynum J, et al: Acute traumatic coagulopathy: the elephant in a room of blind scientists. J Trauma Acute Care Surg 2017;82:S33-S40

Mengoli C, Franchini M, Marano G, et al: The use of fibrinogen concentrate for the management of trauma-related bleeding: a systematic review and meta-analysis. Blood Transfus 2017;15:318-324

Meyer D, Vincent L, Fox E, et al: Every minute counts: time to delivery of initial massive transfusion cooler and its impact on mortality. J Trauma Acute Care Surg 2017;83:19-24

Milton J, Rugino A, Narayan K, et al: A case-based review of the management of penetrating brain trauma. Cureus 2017;9:1342

Min J, Cho H, Jeon S, et al: Effects of 6% hydroxyethyl starch 130/0.4 on postoperative blood loss and kidney injury in off-pump coronary arterial bypass grafting: a retrospective study. Medicine 2017;96:e6801

Mistral T, Boué Y, Bosson J, et al: Performance of point of care international normalized ratio measurement to diagnose trauma-induced coagulopathy. Scand J Trauma Resusc Emerg Med 2017;25:59

Moe D, Lallemand M, McClellan J, et al: Three versus four-factor prothrombin complex concentrates for “factor-based” resuscitation in a porcine hemorrhagic shock model. J Trauma Acute Care Surg 2017;Epub ahead of print

Montgomery H, Butler F, Kerr W, et al: TCCC Guidelines comprehensive review and update: TCCC Guidelines Change 16-03. J Spec Oper Med 2017;17:21-38

Moore H, Moore E, Huebner B, et al: Tranexamic acid is associated with increased mortality in patients with physiological fibrinolysis. J Surg Res 2017;Epub ahead of print

Murray C: Field wound care: prophylactic antibiotics. Wilderness Environ Med 2017;28:S90-S102

National Academies of Science, Engineering, and Medicine. Mil Med 2017;182:1563-1565

Mullally W, Concussion. Am J Med 2017;Epub ahead of print

Neeki M, Dong F, Toy J, et al: Efficacy and safety of tranexamic acid in prehospital traumatic hemorrhagic shock: outcomes of the Cal-PAT study. West J Emerg Med 2017;18:673-683

Nishida T, Kinoshita T, Yamakawa K: Tranexamic acid and trauma-induced coagulopathy. J Intensive Care 2017;5:5

Nutbeam T, Clarke R, Luff T, et al: The height of the cricothyroid membrane of computed tomography scans in trauma patients. Anaesthesia 2017;Epub ahead of print

Oberholzer N, Kaserer A, Albrecht R, et al: Factors influencing quality of pain management in a physician staffed helicopter emergency medical service. Anesth Analg 2017;Epub ahead of print

Ogura T, Lefor A, Nakamura M, et al: Ultrasound-guided resuscitative endovascular balloon occlusion of the aorta in the resuscitation area. J Emerg Med 2017;52:715-722

Ohrt-Nissen S, Bukhari N, Dragsted C, et al: Blood transfusion in the surgical treatment of adolescent idiopathic scoliosis-a single-center experience of patient blood management in 210 cases. Transfusion 2017;Epub ahead of print

Ono Y, Kunil M, Miura T, Shinohara K: "Cannot ventilate, cannot intubate" situation after penetration of the tongue root through to the epipharynx by a surfboard: a case report. J Med Case Rep 2017;11:121

Otten E, Dorlac W: Managing traumatic brain injury: translating military guidelines to the wilderness. Wilderness Environ Med 2017;Epub ahead of print

Parmar P, Greenough P: Optimizing the use of a precious resource: the role of emergency physicians in a humanitarian crisis. West J Emerg Med 2017;18:607-615

Pasarikovski C, Alotaibi N, Al-Mufti F, Macdonald R; Hypertonic saline for raised intracranial pressure following aneurysmal subarachnoid hemorrhage: a systematic review. World Neurosurg 2017;Epub ahead of print.

Patel A, Pieper K, Myburg J, et al: Reanalysis of the Crystalloid versus Hydroxyethyl Starch Trial (CHEST). *N Engl J Med* 2017;377:298-300

Patel E, Ness P, Marshall C, et al: Blood products utilization among trauma and nontrauma massive transfusion protocols at an urban academic medical center. *Anesth Analg* 2017;Epub ahead of print

Patel N, Huang J: Hyperbaric oxygen therapy of spinal cord injury. *Med Gas Res* 2017;7:133-143

Patil P, Chikkapillappa M, Pujara V, et al: ProSeal laryngeal mask airway placement: a comparison of blind versus direct laryngoscopic insertion techniques. *Anesth Essays Res* 2017;11:380-384

Pearce A, Bull A, Clasper J: Mediastinal injury is the strongest predictor of mortality in mounted blast amongst UK deployed forces. *Injury* 2017;Epub ahead of print

Petrosaniak A, Ryzynski A, Lebovic G, Woolfrey K: Cricothyroidotomy in situ simulation curriculum (CRIC Study): training residents for rare procedures. *Simul Healthc* 2017;12:76-82

Phillips J, Mohorn P, Bookstaver R, et al: Hemostatic management of trauma-induced coagulopathy. *Crit Care Nurse* 2017;37:37-47

Pippi R, Santoro M, Cafolla A: The use of chitosan-derived hemostatic agent for postextraction bleeding control in patients on antiplatelet treatment. *J Oral Maxillofac Surg* 2017;75:1118-1123

Plackett T, Cherry D, Delk G, et al: Clinical practice guidelines adherence during Operation Inherent Resolve. *J Trauma Acute Care Surg* 2017;83:S66-S70

Prat N, Meyer A, Ingalls N, et al: ROTEM significantly optimizes transfusion practices for damage control resuscitation in combat casualties. *J Trauma Acute Care Surg* 2017;Epub ahead of print

Press C, Duffy C, Williams J, et al: Measurements of rates of cooling of a manikin insulated with different mountain rescue casualty bags. *Extrem Physiol Med* 2017;Epub ahead of print

Prior S, Cohen M, Conroy A, et al: Correlation between factor (F) Xia, FIXa and tissue factor and trauma severity. *J Trauma Acute Care Surg* 2017;82:1073-1079

Qamarul Hoda M, Samad K, Ullah H: ProSeal versus classic laryngeal mask airway (LMA) for positive pressure ventilation in adults undergoing elective surgery. Cochrane Database Syst Rev 2017;Epub ahead of print

Rall J, Ross J, Clemens M, et al: Hemodynamic effects of the abdominal aortic and junctional tourniquet in a hemorrhagic swine model. J Surg Res 2017;212:159-166

Rand B, Wenke J: An effective negative pressure wound therapy compatible local antibiotic delivery device. J Orthop Trauma 2017;Epub ahead of print

Rasmussen T, Baer D, Remick K, Ludwig G: Combat casualty care research for the multidomain battlefield. J Trauma Acute Care Surg 2017;83:S1-S3

Rasmussen T, Eliason J: Military-civilian partnership in device innovation: development, commercial and application of resuscitative endovascular balloon occlusion of the aorta (REBOA). J Trauma Acute Care Surg 2017; Epub ahead of print

Reith F, Lingsma H, Gabbe B, et al: Differential effects of the Glasgow Coma Scale score and its components: an analysis of 54,069 patients with traumatic brain injury. Injury 2017;Epub ahead of print

Ren Z, Li S, Sheng L, et al: Efficacy and safety of topical use of tranexamic acid in reducing blood loss during primary lumbar spinal surgery: a retrospective case control study. Spine 2017;Epub ahead of print

Reva V, Hörer T, Makhnovskiy A, et al: Field and en route resuscitative endovascular occlusion of the aorta: a feasible military reality? J Trauma Acute Care Surg 2017;83S170-S176

Robak O, Vaida S, Gaitini L, et al: The EasyTube during general anesthesia for minor surgery: a randomized, controlled trial. Medicine 2017;96:e7195

Robak O, Vaida S, Somri M, et al: Inter-center comparison of EasyTube and endotracheal tube during general anesthesia in minor elective surgery. PLoS One 2017;12:e0178756

Roberts H, Osborn P, Wood T, et al: Being prepared for the next conflict: a case analysis of a military Level I trauma center. Mil Med 2017;182:e1681-e1687

Rochlen L, Housey M, Gannon I, et al: Assessing anesthesiology residents' out-of-the-operating room (OOOR) emergent airway management. BMC Anesthesiol 2017;17:96

Rouanet C, Reges D, Rocha E, et al: Traumatic spinal cord injury: current concepts and treatment update. Arq Neuropsiquiatr 2017;75:387-393

Ruetzler K, Guzzella S, Tscholl D, et al: Blind intubation through self-pressurized, disposable supraglottic airway laryngeal intubation masks: an international, multicenter, prospective cohort study. Anesthesiology 2017;Epub ahead of print

Satyarthi G: Biomarker-based targeted therapy of traumatic brain injury: from prehospital care to in-hospital care to rehabilitation. World Neurosurg 2017;103:939-941

Sauter T, Hoess S, Lehmann B, et al: Detection of pneumothoraces in patients with multiple blunt trauma: use and limitations of eFAST. Emerg Med J 2017;Epub ahead of print

Savage S, Zarzaur B, Pohlman T, et al: Clot dynamics and mortality: the MA-R ratio. J Trauma Acute Care Surg 2017;Epub ahead of print

Schauer S, April M, Naylor J, et al: QuikClot® use by ground forces in Afghanistan the prehospital trauma registry experience. J Spec Oper Med 2017;17:101-106

Scherer M, Weightman M, Radomski M, et al: Measuring soldier performance during the patrol-exertion multitask: preliminary validation of a post-concussive functional return to duty metric. Arch Phys Med Rehabil 2017;Epub ahead of print

Seddighi A, Nikouei A, Seddighi A, et al: The role of tranexamic acid in prevention of hemorrhage in major spinal surgeries. Asian J Neurosurg 2017;12:501-505

Seto S, Itakura A, Okagaki R, et al: An algorithm for the management of coagulopathy in postpartum hemorrhage, using fibrinogen concentrate as first-line therapy. Int J Obstet Anesth 2017;Epub ahead of print

Shah S, Spinella P, Muszynski J: Immunologic effects of trauma and transfusion. J Trauma Acute Care Surg 2017;82:S50-S56

Shahid M, Kundra R: Platelet-rich plasma (PRP) for knee disorders. EFFORT Open Rev 2017;2:28-34

Shavit I, Aviram E, Hoffmann Y, et al: Laryngeal mask airway as a rescue device for failed endotracheal intubation during scene-to-hospital air transport of combat casualties. Eur J Emerg Med 2017;Epub ahead of print

Shinkawa T, Holloway J, Tang X, et al: Experience using kaolin-impregnated sponge to minimize perioperative bleeding in Norwood operation. World J Pediatr Congenit Heart Surg 2017;8:475-479

Shlaifer A, Yitzhak A, Baruch E, et al: Point of injury tourniquet application during Operation Protective Edge - what do we learn? Mil Med 2017;83:278-283

Sigal A, Martin A, Ong A: Availability and use of hemostatic agents in prehospital trauma patients in Pennsylvania translation from the military to the civilian setting. Open Access Emerg Med 2017;9:47-52

Sin B, Tatunchak T, Paryavi M, et al; The use of ketamine for acute pain treatment of pain: a randomized, double-blind, placebo-controlled trial. J Emerg Med 2017;52:601-608

Singh K, Gurha P: Comparative evaluation of Ambi AuraGain™ with ProSeal™ laryngeal mask airway in patients undergoing laparoscopic cholecystectomy. Indian J Anaesth 2017;61:496-474

Singleton W, McLean M, Smale M, et al: An analysis of the temperature in change in warmed intravenous fluids during administration in cold environments. Air Med J 2017;36:127-130

Smith W: Integration of tactical EMS in the National Park Services. Wilderness Environ Med 2017;Epub ahead of print

Smith W, Ghazarian N, Christian G: Acute and hyper-acute thoracolumbar corpectomy for traumatic burst fracture using a mini-open lateral approach. Spine 2017;Epub ahead of print

Sobiech P, Adamiak Z, Holak P, et al: Thromboelastographic profile of goat blood after the experimental injury of the femoral artery and use of QuikClot gauze and Celox gauze dressings. Pol J Vet Sci 2017;20:45-49

Söderlund T, Ketonen T, Handolin L: Bleeding pelvic fracture patients: evolution of resuscitation protocols. Scand J Surg 2017;Epub ahead of print

Spaite D, Hu C, Bobrow B, et al: Association of out-of-hospital hypotension depth and duration with traumatic brain injury. Ann Emerg Med 2017;Epub ahead of print

Spinella P: Zero preventable deaths after traumatic brain injury: an achievable goal. *J Trauma Acute Care Surg* 2017;82:S2-S8

Stein P, Kaserer A, Sprengel K, et al: Change of transfusion and treatment paradigm in major trauma patients. *Anaesthesia* 2017;Epub ahead of print

Stevens W, Morse B, Bernard A, et al: Incompatible type A plasma transfusion in patients requiring massive transfusion protocol: outcomes of an Eastern Association for the Surgery of Trauma multicenter study. *J Trauma Acute Care Surg* 2017;83:25-2

Studer N, April M, Bowling F, et al: Albumin for prehospital fluid resuscitation of hemorrhagic shock in tactical combat casualty care. *J Spec Oper Med* 2017;17:82-88

Tanizaki S, Maeda S, Sera M, et al: Small tube thoracostomy (20-22 Fr) in emergent management of chest trauma. *Injury* 2017;Epub ahead of print

Tatum J, Melo N, Ko A, et al: Validation of a field spinal motion restriction protocol in a level I trauma center. *J Surg Res* 2017;Epub ahead of print.

Taylor J, Harvin J, Martin C, et al: Vascular complications from resuscitative endovascular balloon occlusion of the aorta: life over limb? *J Trauma Acute Care Surg* 2017;83:S120-S123

te Grotenhuis R, van Grunsven P, Heutz W, Tan E: Prehospital use of hemostatic dressings in emergency medical services in the Netherlands: a prospective study of 66 cases. *Injury* 2016

Thitivaraporn P, Narueponjirakul N, Samorn P, et al: Randomized controlled trial of chest tube removal aided by a party balloon. *Asian Cardiovasc Thorac Ann* 2017;Epub ahead of print

Thompson S, Wilson Z, Cole C, et al: Case report: mass casualty lightning strike at Ranger Training Camp. *Mil Med* 2017;182:e1803-e1806

Tien A, Beckett A, Pannell D: The University of Toronto's lasting contribution to war surgery: how Maj. L. Bruce Robertson fundamentally transformed thinking toward blood transfusion during the First World War. *Can J Surg* 2017;60:152-154

Trimmel H, Majdan M, Wodak A, et al: Citicoline in severe traumatic brain injury: indications for improved outcomes: a retrospective matched pair analysis from 14 Austrian trauma centers. *Wien Klin Wochenschr* 2017;Epub ahead of print

Turco L, Cornell D, Phillips B: Penetrating bihemispheric traumatic brain injury: a collective review of gunshot wounds to the head. *World Neurosurg* 2017;Epub ahead of print

Turner C, Stockinger Z, Gurney J: Combat surgical workload in Operation Iraqi Freedom and Operation Enduring Freedom: the definitive analysis. *J Trauma Acute Care Surg* 2017;83:77-83

Uchino H, Kong V, Oosthuizen G, et al: Can we train military surgeons in a civilian trauma center? *World J Surg* 2017

van Rein E, Houwert, Gunning A, et al: Accuracy of prehospital triage protocols in selecting severely injured patients: a systematic review. *J Trauma Acute Care Surg* 2017;83:328-339

van Zundert A, Gatt S, Gatt S, et al: 'Failed supraglottic airway': an algorithm for suboptimally placed supraglottic airway devices based on videolaryngoscopy. *Br J Anaesth* 2017;118:645-649

Vertu N, Nascimento M, Pasquier P: Buddy-aid battlefield pain management. *J Trauma Acute Care Surg* 2017;Epub ahead of print

Vithalani V, Vil S, David S, Richmond N: Unrecognized failed airway management using a supraglottic airway device. *Resuscitation* 2017;Epub ahead of print

Wallace M, Britton S, Meek R, et al: Comparison of five video-assisted intubation devices by novice and expert laryngoscopists for use in the aeromedical evacuation environment. *Mil Med Res* 2017;4:20

Watermeyer M, Dippenaar N, Simo N, et al: Essential lessons in a potential sarin attack disaster plan for a resource-constrained environment. *Disaster Med Public Health Prep* 2017;Epub ahead of print

Wedmore I, Butler F: Battlefield analgesia in tactical combat casualty care. *Wilderness Environ Med* 2017;28:S109-S116

White N, Ward K, Pati S, et al: Hemorrhagic blood failure: oxygen debt, coagulopathy, and endothelial damage. *J Trauma Acute Care Surg* 2017;82:S41-S56

Wilson S, Gangathimmaih V: Does prehospital management by doctors affect outcomes in major trauma? A systematic review. *J Trauma Acute Care Surg* 2017;Epub ahead of print

Winearls J, Wullscheher M, Wake E, et al: Fibrinogen early in severe trauma study (FIELTY): study protocol for a randomised controlled trial. *Trials* 2017;18:241

Wurster L, Thakkar R, Haley K, et al: Standardizing the initial resuscitation of the trauma patient with the primary assessment completion tool using video review. *J Trauma Acute Surg Care* 2017;82:1002-1006

Xi C, Zhu L, Zhuang Y, et al: Experimental evaluation of tranexamic acid-loaded porous starch as a hemostatic powder. *Clin Appl Thromb Hemost* 2017;Epub ahead of print

Xu R, Zhu Y, Fan Q, et al: Comparison between the cobra perilaryngeal airway management and laryngeal mask airways under general anesthesia: a systematic review and meta-analysis. *Anesth Analg* 2017;Epub ahead of print

Yang L, Du S, Sun Y: Is combined topical and intravenous tranexamic acid superior to single use of tranexamic acid in total joint arthroplasty?: A meta-analysis from randomized trials. *Medicine* 2017;96:e7609

Yitzhak A, Glick Y, Benov A, et al: The need for optimized crystalloid-based resuscitation. *J Trauma Acute Care Surg* 2017;82:S66-S69

Yonemura S, Doane S, Keil S, et al: Improving the safety of whole blood-derived transfusion products with riboflavin-based pathogen reduction technology. *Blood Transfus* 2017;15:357-364

Zhang H, He G, Zhang C, et al: Is combined topical and intravenous tranexamic acid superior to intravenous tranexamic acid alone for controlling blood loss after total hip arthroplasty?: A meta-analysis. *Medicine* 2017;96:e6916

Zhao W, Chen B, Wang J, et al: Establishment of an ideal time window model in hypothermic-targeted temperature management after traumatic brain injury in rats. *Brain Res* 2017;Epub ahead of print

Zielinski M, Stubbs J, Berns K, et al: Prehospital blood transfusion programs: capabilities and lessons learned. *J Trauma Acute Care Surg* 2017;82:S70-S78

Zufferey P, Lanoiselée J, Chapelle C, et al: Intravenous tranexamic acid bolus plus infusion is not more effective than a single bolus in primary hip arthroplasty: a randomized controlled trial. Anesthesiology 2017;Epub ahead of print